

Drug Checking in Interior Health

Key Findings

- The percentage of opioids testing positive for benzodiazepines (22.9%) in the Interior Health region has continued on an upward trend since September.
- No alerts were issued for samples checked at FTIR/test strip checking services this month.










Drug Checking in Interior Health


Number of samples tested with fentanyl present

















PUBLIC HEALTH NOTIFICATIONS

Date & Location	Expected	Drugs Detected	Fentanyl	Benzo	Area	Alert Message
	Drug		Strip	Strip	Purchased	

No alerts from Interior Health partners this December

For information about public health alerts in the Interior Health region, please visit:



During the month of December, 22.9% of expected opioid samples tested positive for benzodiazepines using test strips in our partner sites in the Interior Health region (8 samples of 35 tested).

Opioid samples are tested for benzodiazepine-positivity using BTNX test strips. The results presented above are presumptive and tentative until confirmed by a laboratory.

Number of drugs checks at each site

Location	# of Drugs Checked		
ANKORS (Nelson)	43		
ASK Wellness Head Office (Kamloops)	11		
ASK Wellness Burdock House (Penticton)	3		
ASK Wellness (Merritt)	9		
Interior Health (Kelowna)	8		
Interior Health (Vernon)	5		
Total	79		

Number of samples that matched expectation

using FTIR/test strip drug checking

EXPECTED DRUG:

Depressant

2 Samples Tested



Matched: 2

EXPECTED DRUG:

5 Samples Tested



Match not determined: 5

Psychedelic

17 Samples Tested



Matched: 14 Did not match: 3 EXPECTED DRUG:

Stimulant

17 Samples Tested



Matched: 17

EXPECTED DRUG: Polysubstance

1 Sample Tested



Matched: 1

EXPECTED DRUG:

Opioid

37 Samples Tested



Matched: 36 Did not match: 1

Total

79 Samples Tested



Matched: 70 Did not match: 4

Match not determined: 5

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

Drug Checking in Interior Health

Number of opioid samples that matched expectation

using FTIR/test strip drug checking

EXPECTED DRUG:

Fentanyl

35 Samples Tested



EXPECTED DRUG:

Heroin

2 Samples Tested



Matched: 1

Did not match: 1

Matched: 35

Total

37 Samples Tested



Matched: 36 Did not match: 1

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents. 'Down' can refer to any opioid drug present in any amount.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazapine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of both the FTIR and BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners:









