
Drug Checking in Fraser Health


Key Findings

- Low numbers of samples are a result of SafePoint drug checking services moving to a drop-off model due to pandemic restrictions. Samples can be dropped-off during operating hours (7am-1am) and results are available after Tuesday morning each week.
- One sample of expected fentanyl contained a synthetic cannabinoid, AMB-FUBINACA.
- Four Xanax (alprazolam) tablets were submitted for checking this month. One was found to contain etizolam, one
 contained alprazolam, and the remaining two tested positive for benzodiazepines (undetermined).



Components found in drug samples in March


1 sample of expected fentanyl contained Caffeine, a sugar alcohol, and AMB-FUBINACA, a synthetic cannabinoid.





Drug Checking in Fraser Health

Number of samples tested with fentanyl present



0%

OCTOBER

1 Stimulant samples



NOVEMBER

4 Depressant samples
4 FENTANYL NEGATIVE (100%)



2 Psychedelic samples



FEBRUARY

MARCH

1 Unknown samples
1 FENTANYL NEGATIVE (100%)






PUBLIC HEALTH NOTIFICATIONS

| Date & Location | Expected Drug | Drugs Detected | Fentanyl Strip | Benzo Strip | Alert Message | | |
|--|------------------|----------------|-------------------|----------------|---------------|--|--|
| No alerts from Fraser Health partners this January | | | | | | | |

For information about current substances in the Fraser Health region: overdose@fraserhealth.ca

Percentage of opioids testing positive for benzodiazepines in the past 6 months



Month

JANUARY

DECEMBER

During the month of March, **0**% of expected opioid samples tested positive for benzodiazepines using test strips in our partner sites in the Fraser Health Region (**0** samples of **3 tested**). *These numbers should be taken with caution as this trend is based on low samples.

Opioid samples are tested for benzodiazepine-positivity using BTNX test strips. The results presented above are presumptive and tentative until confirmed by a laboratory.

Drug Checking in Fraser Health

Number of drugs checks at each site

| Location | # of drug checks | Hours | |
|--------------------|------------------|--|--|
| SafePoint (Surrey) | 11 | SafePoint is currently accepting samples for checking during operating hours. A technician checks samples every Tuesday morning and results are available after 12PM. | |
| | | For details visit https://drugcheckingbc.ca/drug-checking-sites/fraser-health/ | |

of samples that matched client expectation

using FTIR/test strip drug checking

EXPECTED DRUG: Opioid

3 Samples Tested

Matched: 3

Stimulant

1 Samples Tested

•

Matched: 1

EXPECTED DRUG:

Depressant

4 Samples Tested

Matched: 3
Did not match: 1

EXPECTED DRUG:

Psychedelic

2 Samples Tested

Matched: 2

EXPECTED DRUG:

Unknown

1 Samples Tested

Matched not determined: 1

Total

11 Samples Tested



Matched: 9

Did not match: 1

Matched not determined: 1

Number of opioid samples that matched client expectation

using FTIR/test strip drug checking

EXPECTED DRUG: Fentanyl

2 Samples Tested



Matched: 2

EXPECTED DRUG:

Pharmaceutical

1 Samples Tested



Matched: 1

Total

3 Samples Tested



Matched: 3

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

Depressant may include: benzodiazepines, etizolam, GHB, hypnotics Opioid may include: "down," heroin, fentanyl, pharmaceutical opioids

Polysubstance includes: cross-category mixtures

Psychedelic may include: MDMA and related, 2C-family, tryptamines, ketamine. LSD

Stimulant may include:

methamphetamine, "speed," cocaine and crack cocaine, cathinones Unknown includes:

Samples where the individual was unable to identify an expected substance. This includes found samples.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazapine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of both the FTIR and BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners:







