


## Key Findings


- The percentage of opioids testing positive for benzodiazepines rose to 46% (12 of 26 samples checked). Trends may be hard to infer due to the small number of samples. Etizolam, the predominant benzodiazepine in expected opioids, may be missed by drug checking technologies. This means that the true rate may be higher than reported here.
- Drug checking services are available Thursdays and Fridays from 11:30am–3:30pm at 437 Wesley Street. Samples can also be dropped off any day of the week and results can be picked up during drug checking hours.
- The Vancouver Island Drug Checking Project offers drug checking in Victoria. For information about their project and more drug checking sample results, please visit [substance.uvic.ca](http://substance.uvic.ca).

## Number of drug samples checked in the past 6 months




## Components found in drug samples in September

### Components in opioids



## Number of samples tested with fentanyl present




## PUBLIC HEALTH NOTIFICATIONS

| Date & Location                        | Expected Drug | Drugs Detected | Fentanyl Strip | Benzo Strip | Alert Message |
|--|---------------|----------------|----------------|-------------|---------------|
| No alerts for Island Health this month |               |                |                |             |               |

For information about current substances in the Island Health region: <https://substance.uvic.ca/>

## Percentage of opioids testing positive for benzodiazepines in the past 6 months








During the month of September, **46.2%** of expected opioid samples tested positive for benzodiazepines using test strips in our partner site in Island Health (**12 samples of 26 checked**). Opioid samples are checked for benzodiazepine-positivity using BTNX test strips and the FTIR spectrometer. The results presented here are derived from both of these technologies and are presumptive until confirmed by a laboratory.

## Number of drug checks at each site:

| Location                 | # of Drug Checks |
|--------------------------|------------------|
| Wesley Street OPS (CMHA) | 32               |

## Number of samples that matched client expectation

using FTIR/test strip drug checking

| EXPECTED DRUG:   | EXPECTED DRUG:  | EXPECTED DRUG:  | EXPECTED DRUG:   | EXPECTED DRUG:  |
|--|---|---|--|---|
| <b>Opioid</b><br>26 Samples Tested   | <b>Stimulant</b><br>1 Samples Tested  | <b>Depressant</b><br>2 Samples Tested   | <b>Psychedelic</b><br>2 Samples Tested   | <b>Unknown</b><br>1 Samples Tested  |
|  |  |  |  |  |
| Matched: 23<br>Did not match: 3  | Matched: 1  | Matched: 1<br>Did not match: 1  | Matched: 2   | Match not determined: 1   |

### Total




32 Samples Tested



Matched: 27  
Did not match: 4  
Match not determined: 1

## Number of opioid samples that matched client expectation

\*using FTIR/test strip drug checking

| EXPECTED DRUG:   | EXPECTED DRUG:  | EXPECTED DRUG:  |
|--|---|---|
| <b>Down</b><br>15 Samples Tested   | <b>Heroin</b><br>3 Sample Tested  | <b>Fentanyl</b><br>8 Sample Tested  |
|  |  |  |
| Matched: 15  | Did not match: 3  | Matched: 8  |

### Total

26 Samples Tested



Matched: 23  
Did not match: 3

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

**Depressant may include:**  
benzodiazepines, etizolam,  
GHB, hypnotics

**Opioid may include:**  
"down," heroin, fentanyl,  
pharmaceutical opioids

**Polysubstance includes:**  
cross-category mixtures

**Psychedelic may include:**  
MDMA and related,  
2C-family, tryptamines,  
ketamine, LSD

**Stimulant may include:**  
methamphetamine,  
"speed," cocaine and crack  
cocaine, cathinones

**Unknown includes:**  
samples where the individual  
was unable to identify an  
expected substance - this  
includes found samples.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazepine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of both the FTIR and BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners: