

# **Drug Checking in Island Health**

#### **Key Findings**

- The percentage of opioids testing positive for benzodiazepines in the region was 76.9% (10 of 13 samples). Trends may be hard to infer due to the small number of samples over a large region. The true rate may be higher than reported here because etizolam, the predominant benzodiazepine in expected opioids, may be missed by drug checking technologies.
- Drug checking services are available Thursdays and Fridays from 11:30am-3:30pm at 437 Wesley Street. Samples can also be dropped off any day of the week and results can be picked up during drug checking hours.
- The Vancouver Island Drug Checking Project offers drug checking in Victoria. For information about their project and more drug checking sample results, please visit substance.uvic.ca.








# **Drug Checking in Island Health**

## Number of samples tested with fentanyl present






#### **PUBLIC HEALTH NOTIFICATIONS**

Date & Location	Expected Drug	Drugs Detected	Fentanyl Strip	Benzo Strip	Alert Message
-----------------	------------------	----------------	-------------------	----------------	---------------

No alerts for Island Health this month

For information about current substances in the Island Health region: https://substance.uvic.ca/

#### Percentage of opioids testing positive for benzodiazepines in the past 6 months



During the month of May, **76.9**% of expected opioid samples tested positive for benzodiazepines using test strips in our partner site in Island Health **(10 samples of 13 checked)**. Opioid samples are checked for benzodiazepine-positivity using BTNX test strips and the FTIR spectrometer. The results presented here are derived from both of these technologies and are presumptive until confirmed by a laboratory.

# **Drug Checking in Island Health**

## Number of drugs checks at each site:

Location	# of Drug Checks		
Wesley Street OPS (CMHA)	20		

# Number of samples that matched client expectation

using FTIR/test strip drug checking

**Opioid** 

13 Samples Tested

Matched: 13

**EXPECTED DRUG:** 

**Stimulant** 

4 Samples Tested

Matched: 4

**EXPECTED DRUG:** 

**Depressant** 

1 Samples Tested

Did not match: 1

EXPECTED DRUG:

**Psychedelic** 

2 Samples Tested

Matched: 2

**Total** 

20 Samples Tested

Matched: 19

Did not match: 1

## Number of opioid samples that matched client expectation

\*using FTIR/test strip drug checking

**EXPECTED DRUG:** 

### **Fentanyl**

1 Samples Tested

Matched: 1

**EXPECTED DRUG:** 

Heroin

1 Samples Tested

ĺ

Matched: 1

**EXPECTED DRUG:** 

Down

10 Samples Tested

Matched: 10

**EXPECTED DRUG:** 

**Pharmaceutical** 

1 Samples Tested

Matched: 1

**Total** 

13 Samples Tested

Matched: 13

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

**Depressant may include:** benzodiazepines, etizolam, GHB, hypnotics **Opioid may include:** "down," heroin, fentanyl, pharmaceutical opioids

Polysubstance includes: cross-category mixtures

Psychedelic may include:

MDMA and related,

2C-family, tryptamines,

ketamine, LSD

Stimulant may include: methamphetamine, "speed," cocaine and crack cocaine, cathinones Unknown includes: samples where the individual was unable to identify an expected substance - this

includes found samples.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazapine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of <u>both</u> the FTIR <u>and</u> BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners:





