

Drug Checking in Northern Health

Key Findings

- Drug checking services with FTIR and test strips are now available in the Northern Health Region. Services are available at POUNDS in Prince George and the Northwest Intensive Case Management Team (ICMT) location in Terrace. For updated times and locations, visit https://www.northernhealth.ca/health-topics/overdose-prevention#drug-checking
- A total of 49 drug checks were conducted across these two sites in the Northern Health region in August, up from 26 samples the previous month. Opioids and stimulants comprised the majority of samples checked.
- The percentage of opioids testing positive for benzodiazepines in the region increased from 37.5% to 44.8% in August (13 of 29 samples). The true rate may be higher than reported here as etizolam, a benzodiazepine-like substance, may be missed by drug checking technologies.
- From the 49 samples tested in August, results seem consistent with other areas of the province. Fentanyl was detected in 28 out of 29 opioid samples checked. 1 stimulant sample tested positive for fentanyl in August.






During the month of August, **44.8**% of expected opioid samples tested positive for benzodiazepines using test strips in our partner sites in Northern Health **(13 samples of 29 checked)**. Opioid samples are checked for benzodiazepine-positivity using BTNX test strips and the FTIR spectrometer. The results presented here are derived from both of these technologies and are presumptive until confirmed by a laboratory.





Drug Checking in Northern Health

Number of samples tested with fentanyl present












1 Other



1 Polysubstance





PUBLIC HEALTH NOTIFICATIONS

Date & Location	Expected Drug	Drugs Detected	Fentanyl Strip	Benzo Strip	Alert Message

No alerts from Northern Health partners this month

Number of drugs checks at each site

Location	# of drug checks		
POUNDS (Prince George)	32		
Northwest ICMT (Terrace)	14		
Overdose Prevention Community Health Fair	3		

Drug Checking in Northern Health

of samples that matched client expectation

using FTIR/test strip drug checking

EXPECTED DRUG:

Opioid

29 Samples Tested



Matched: 28

Did Not Match: 1

EXPECTED DRUG: Other

1 Samples Tested

Matched: 1

EXPECTED DRUG:

Stimulant

12 Samples Tested



Matched: 11

EXPECTED DRUG:

Match not determined: 1

Polysubstance

1 Samples Tested

Matched: 1

EXPECTED DRUG:

Depressant

1 Samples Tested

EXPECTED DRUG: Unknown

2 Samples Tested



Did Not Match: 1

Match not determined: 2

Total

49 Samples Tested



Matched: 43

Did Not Match: 3 Match not determined: 3

Number of opioid samples that matched client expectation

using FTIR/test strip drug checking

EXPECTED DRUG:

Fentanyl

1 Samples Tested

Matched: 1

EXPECTED DRUG:

Down

28 Samples Tested



Matched: 27 Did Not Match: 1 **EXPECTED DRUG:**

Total

29 Samples Tested



EXPECTED DRUG:

Matched: 2

Did Not Match: 1

Psychedelic

3 Samples Tested

Matched: 28 Did Not Match: 1

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

Depressant August include: benzodiazepines. etizolam. GHB, hypnotics

Opioid August include: "down," heroin, fentanyl, pharmaceutical opioids

Polysubstance includes: cross-category mixtures

Psychedelic August include: MDMA and related, 2C-family, tryptamines, ketamine, LSD Stimulant August include: methamphetamine. "speed," cocaine and crack cocaine, cathinones

Unknown includes:

Samples where the individual was unable to identify an expected substance. This includes found samples.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazapine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of both the FTIR and BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners:





