

Drug Checking in Island Health

Key Findings

- In March, a total of 182 drug checks were performed at the Albert Street site, nearly doubling last month's record high. Increases in samples checked occurred primarily in the opioid category.
- The percentage of opioids testing positive for benzodiazepines decreased from 88.7% to 61.2% in March (90 of 147 samples).
- Drug checking services are expanding! Services are now operating 7 days a week from 11:00am-8:30pm at 250 Albert Street. Samples can also be dropped off any day of the week and results can be picked up during drug checking hours.
- Substance (The Vancouver Island Drug Checking Project), offers drug checking in Victoria and communities across the Island, as well as mail in services. For more information about drug checking sample results in the Island health region, please visit substance.uvic.ca.











Drug Checking in Island Health


Number of samples tested with fentanyl present













PUBLIC HEALTH NOTIFICATIONS

Date & Location

Expected Drug

Drugs Detected

Fentanyl Strip


Benzo Strip

Alert Message

No alerts from drugs checked at Wesley St Drug Checking service this month

For information about current substances in the Island Health region: https://substance.uvic.ca/

Percentage of opioids testing positive for benzodiazepines in the past 6 months



During the month of March, **61.2**% of expected opioid samples tested positive for benzodiazepines using test strips in our partner site in Island Health **(90 samples of 147 checked)**. Opioid samples are checked for benzodiazepine-positivity using BTNX test strips and the FTIR spectrometer. The results presented here are derived from both of these technologies and are presumptive until confirmed by a laboratory.


Drug Checking in Island Health

Number of drugs checks at each site:

Location	# of Drug Checks
Wesley Street OPS (CMHA)	182

Number of samples that matched client expectation

using FTIR/test strip drug checking



Number of opioid samples that matched client expectation

*using FTIR/test strip drug checking



Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

Depressant include: benzodiazepines, etizolam, GHB, hypnotics **Opioid include:**"down," heroin, fentanyl,
pharmaceutical opioids

Polysubstance includes: cross-category mixtures

Psychedelic include: MDMA and related, 2C-family, tryptamines, ketamine, LSD

Stimulant include: methamphetamine, "speed," cocaine and crack cocaine, cathingnes Unknown includes: samples where the individual was unable to identify an expected substance - this includes found samples.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazapine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of <u>both</u> the FTIR <u>and</u> BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners:















