

Key Findings

- In May, a total of 286 drug checks were performed by services offering FTIR drug checking in the Interior Health region.
- The percentage of opioids testing positive for benzodiazepines increased from 73.6% to 88.8% in May (142 of 160 samples). Trends may be hard to infer due to the large geographic region.
- Four drug alerts were issued by Interior Health, three pertaining to down samples that contained high concentrations of fentanyl and bromazolam. One alert concerned a methamphetamine sample that instead tested as fentanyl.
- Drug checking is available in several communities across the Interior Health region. For updated times and locations, visit https://drugchecking.ca/







Number of samples tested with fentanyl present

160 Opioid samples

4 FENTANYL NEGATIVE (2%)



156 FENTANYL POSITIVE

63 Stimulant samples

62 FENTANYL NEGATIVE (98%)



1 FENTANYL POSITIVE

5 Depressant samples

2 FENTANYL NEGATIVE



1 FENTANYL 2 FENTANYL NO POSITIVE (20%) TESTED (40%)

33 Psychedelic samples

33 FENTANYL NEGATIVE (100%)



20 Unknown samples

12 FENTANYL NEGATIVE (60%)



8 FENTANYL POSITIVE

5 Polysubstance samples

5 FENTANYL POSITIVE (100%)





PUBLIC HEALTH NOTIFICATIONS

Date & Location	Expected Drug	Drugs Detected	Fentanyl Strip	Benzo Strip	Area Purchased	Alert Message
May 4, 2023 Crankbrook	Down	Caffeine, Erythritol, Fentanyl, Bromazolam	Positive	Positive	Cranbrook	A substance sold as down tested positive for fentanyl, fentanyl analogues, and bromazolam in Cranbrook.
May 12, 2023 Penticton	Down	Fentanyl, Bromazolam, Caffeine	Positive	Positive	Penticton	Light blue chunks sold as down tested positive for fentanyl and bromazolam in Penticton.
May 25, 2023 Kelowna	Down	Caffeine, Fentanyl, Bromazolam	Positive	Positive	Kelowna	Black chunks sold as down tested positive for fentanyl and bromazolam in Kelowna.
May 29, 2023 Penticton	Methamphetamine	Fentanyl	Positive	Negative	Penticton	White crystals and powder sold as methamphetamine tested positive for fentanyl in Penticton.

Health authorities and community organizations issue further toxic drug alerts from sources other than drug checking.

See their respective websites or social media accounts for more alerts.

For information about public health alerts in the Interior Health region, please visit:

Percentage of opioids testing positive for benzodiazepines in the past 6 months



During the month of May, **88.8**% of expected opioid samples tested positive for benzodiazepines using test strips in our partner sites in Interior Health **(142 samples of 160 checked)**. Opioid samples are checked for benzodiazepine-positivity using BTNX test strips and the FTIR spectrometer. The results presented here are derived from both of these technologies and are presumptive until confirmed by a laboratory.


Number of drugs checks at each site

Location	# of Drugs Checked		
ANKORS (Cranbrook)	70		
ANKORS (Grand Forks)	25		
ANKORS (Nelson)	30		
ANKORS (Trail)	21		
ASK Wellness (Kamloops)	12		
ASK Wellness (Penticton)	74		
UBCO HaRT (Kelowna)	50		
Total	286		

Less than 5 samples were tested at the following location, which is included in the total: ANKORS (Kimberley).

Number of samples that matched expectation

using FTIR/test strip drug checking




Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

includes found samples.

Number of opioid samples that matched expectation

using FTIR/test strip drug checking



Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents. 'Down' can refer to any opioid drug present in any amount.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazapine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of both the FTIR and BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners:









