


Key Findings

- In July, a total of 194 drug checks were performed at the Albert Street harm reduction services site, the highest number of samples recorded by the site to date. Increases occurred primarily in the opioid and stimulant categories.
- The percentage of opioids testing positive for benzodiazepines by FTIR and test strip increased from 73.3% to 91.1% in July (112 of 123 samples).
- Bromazolam, a benzodiazepine, was found in concentrations high enough to be detected by FTIR in 36.0% of unregulated opioid samples checked (27 of 75 samples).
- Drug checking services are expanding! Services are now operating 7 days a week from 11:00am–8:30pm at 250 Albert Street. Samples can also be dropped off any day of the week and results can be picked up during drug checking hours.
- Substance (The Vancouver Island Drug Checking Project), offers drug checking in Victoria and communities across the Island, as well as mail in services. For more information about drug checking sample results in the Island health region, please visit substance.uvic.ca.

Number of drug samples checked in the past 6 months




Number of samples by drug category in July

Opioid	123
Stimulant	57
Depressant	2
Psychedelic	5
Other	0
Polysubstance	0
Unknown	7

TOTAL: 194


Components found in drug samples in July

Components in unregulated opioids




Additional compounds that were detected a single time each have been excluded from this graph, which include: cocaine base, cocaine hcl, heroin, mdma, quetiapine.

Components in cocaine



Number of samples tested with fentanyl present




PUBLIC HEALTH NOTIFICATIONS

Date & Location	Expected Drug	Drugs Detected	Fentanyl Strip	Benzo Strip	Alert Message
No alerts from drugs checked at Albert St. Drug Checking service this month					

For information about current substances in the Island Health region: <https://substance.uvic.ca/>

Percentage of opioids testing positive for benzodiazepines in the past 6 months



During the month of July, **91.1%** of expected opioid samples tested positive for benzodiazepines using test strips in our partner site in Island Health (**112 samples of 123 checked**). Opioid samples are checked for benzodiazepine-positivity using BTNX test strips and the FTIR spectrometer. The results presented here are derived from both of these technologies and are presumptive until confirmed by a laboratory.

Number of drugs checks at each site:

Location	# of Drug Checks
Albert St. OPS (CMHA Nanaimo)	182
Laketown Shakedown Music Festival	12
Total:	194

Number of samples that matched client expectation

using FTIR/test strip drug checking

EXPECTED DRUG	Matched	Did Not Match	Match Not Determined	Samples Tested
Opioid	118	5	0	123
Stimulant	52	5	0	57
Depressant	2	0	0	2
Psychedelic	4	1	0	5
Unknown	0	0	7	7
Total	176	11	7	194

Number of opioid samples that matched client expectation

*using FTIR/test strip drug checking

EXPECTED DRUG	Matched	Did Not Match	Match Not Determined	Samples Tested
Fentanyl	4	0	0	4
Other Opioid	0	1	0	1
Down	114	3	0	117
Pharmaceutical	0	1	0	1
Total	118	5	0	123

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

Depressant include:
benzodiazepines, etizolam,
GHB, hypnotics

Opioid include:
"down", heroin, fentanyl
(unregulated opioids), and
pharmaceutical opioids

Polysubstance includes:
cross-category mixtures

Psychedelic include:
MDMA and related,
2C-family, tryptamines,
ketamine, LSD

Stimulant include:
methamphetamine, "speed,"
cocaine and crack cocaine,
cathinones

Unknown includes:
samples where the individual
was unable to identify an
expected substance - this
includes found samples.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazepine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of both the FTIR and BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners: