


Key Findings

- In September, a total of 48 drug checks were performed by services offering FTIR drug checking in the Northern Health region. This increase was due to POUNDS resuming normal operations in September.
- The percentage of opioids testing positive for benzodiazepines by FTIR and test strip decreased from 100% to 38.1% in September (8 of 21 samples), with bromazolam most frequently detected.
- Fluorofentanyl was detected in 15.0% of unregulated opioid samples (3 of 20 samples), whereas fentanyl was detected in 30.0% (6 samples).
- Xylazine was detected in 10.0% of unregulated opioid samples (2 of 20 samples).
- Trends are hard to infer due to the small number of samples over a large region.
- Drug checking services with FTIR and test strips are now available in the Northern Health Region. For updated times and locations, please visit the [Northern Health website](#).

Number of drug samples checked in the past 6 months




Number of samples by drug category in September

Opioid	21
Stimulant	20
Depressant	0
Psychedelic	3
Other	2
Polysubstance	0
Unknown	2


TOTAL: 48

Components found in drug samples in September

Components in unregulated opioids




Components in cocaine



Additional compounds that were detected a single time each have been excluded from this graph, which include: inositol, mannitol, methamphetamine.

Number of samples tested with fentanyl present




PUBLIC HEALTH NOTIFICATIONS

Date & Location	Expected Drug	Drugs Detected	Fentanyl Strip	Benzo Strip	Area Purchased	Alert Message
No alerts for Northern Health this month.						

Number of drugs checks at each site

Location	# of drug checks
POUNDS (Prince George)	34
Terrace ICMT (Terrace)	14
Total	48

Percentage of opioids testing positive for benzodiazepines in the past 6 months



During the month of September, **38.1%** of expected opioid samples tested positive for benzodiazepines using test strips in our partner site in Northern Health (**8 samples of 21 checked**). Opioid samples are checked for benzodiazepine-positivity using BTNX test strips and the FTIR spectrometer. The results presented here are derived from both of these technologies and are presumptive until confirmed by a laboratory.

of samples that matched client expectation using FTIR/test strip drug checking

EXPECTED DRUG	Matched	Did Not Match	Match Not Determined	Samples Tested
Opioid	21	0	0	21
Stimulant	19	1	0	20
Psychedelic	3	0	0	3
Other	2	0	0	2
Unknown	0	0	2	2
Total	45	1	2	48

Number of opioid samples that matched client expectation using FTIR/test strip drug checking

EXPECTED DRUG	Matched	Did Not Match	Match Not Determined	Samples Tested
Fentanyl	1	0	0	1
Down	19	0	0	19
Pharmaceutical	1	0	0	1
Total	21	0	0	21

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

Depressant include:
benzodiazepines, etizolam,
GHB, hypnotics

Opioid include:
"down", heroin, fentanyl
(unregulated opioids), and
pharmaceutical opioids

Polysubstance includes:
cross-category mixtures

Psychedelic include:
MDMA and related,
2C-family, tryptamines,
ketamine, LSD

Stimulant include:
methamphetamine, "speed,"
cocaine and crack cocaine,
cathinones

Unknown includes:
Samples where the individual
was unable to identify an
expected substance. This
includes found samples.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazepine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of both the FTIR and BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners:

