

#### **Key Findings**


- In December, a total of 309 drug checks were performed by services offering FTIR drug checking in the Interior Health region.
- The percentage of all opioids testing positive for benzodiazepines by FTIR and test strip increased from 65.0% to 70.7% in December (99 of 140 samples).
- Benzodiazepines were found in concentrations high enough to be detected by FTIR in 18.4% of unregulated opioids (25 of 136 samples), with bromazolam most frequently detected (24 samples).
- Fluorofentanyl was detected by FTIR in 26.5% of unregulated opioids (36 of 136 samples), whereas fentanyl was detected in 36.0% (49 samples).
- Xylazine was detected by FTIR in 2.9% of unregulated opioid samples (4 of 136 samples).
- Drug checking is available in several communities across the Interior Health region. The Medicine Shoppe Pharmacy
  offers drug checking services in Vernon at #100-3605 31st Street. For updated times and locations in Interior Health,
  visit https://drugchecking.ca/







### **Number of samples tested with fentanyl present**





#### **PUBLIC HEALTH NOTIFICATIONS**


Date & Location	Expected Drug	FTIR Spectroscopy Results	Fentanyl Strip	Benzo Strip	Alert Message
December 18, 2023 Nelson	Down	Caffeine, Fentanyl, Erythritol, Xylazine, Bromazolam	Positive	Positive	Pink chunks sold as down tested positive for high amounts of fentanyl (25%), bromazolam (17%), and xylazine (11%) in Nelson*
December 21, 2023 Cranbrook	Down	Caffeine, Fentanyl, Erythritol, Xylazine, Uncertain Match	Positive	Positive	Pink chunks sold as down tested positive for high amounts of fentanyl (27%), bromazolam (17%), and xylazine (26%) in Cranbrook*

\*Quantitative results provided by complementary testing partner Substance using Paper Spray Mass Spectrometry, all other spectroscopy results are determined by FTIR.

Health authorities and community organizations issue further toxic drug alerts from sources other than drug checking.

See their respective websites or social media accounts for more alerts.

For information about public health alerts in the Interior Health region, please visit:




During the month of December, **70.7**% of expected opioid samples tested positive for benzodiazepines using test strips in our partner sites in Interior Health **(99 samples of 140 checked)**. Opioid samples are checked for benzodiazepine-positivity using BTNX test strips and the FTIR spectrometer. The results presented here are derived from both of these technologies and are presumptive until confirmed by a laboratory.

### Number of drugs checks at each site

Location	# of Drugs Checked
ANKORS (Cranbrook)	45
ANKORS (Grand Forks)	17
ANKORS (Nelson)	64
ANKORS (Trail)	15
ASK Wellness (Kamloops)	25
ASK Wellness (Penticton)	30
Medicine Shoppe Pharmacy (Vernon)	12
UBCO HaRT (Kelowna)	101
Total:	309

### Number of samples that matched expectation

using FTIR/test strip drug checking




Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

includes found samples.

### Number of opioid samples that matched expectation

using FTIR/test strip drug checking



Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents. 'Down' can refer to any opioid drug present in any amount.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazapine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of both the FTIR and BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

#### BCCSU gratefully acknowledges the contributions of the following partners:









