


February 2024

Key Findings

- In February, a total of 430 drug checks were performed by services offering FTIR drug checking in the Interior Health region.
- The percentage of all opioids testing positive for benzodiazepines by FTIR and test strip decreased from 76.0% to 73.3% in February (137 of 187 samples).
- Benzodiazepines were found in concentrations high enough to be detected by FTIR in 29.3% of unregulated opioids (51 of 174 samples), with bromazolam most frequently detected (29 samples), followed by desalkylgidazepam (19 samples).
- Fluorofentanyl was detected by FTIR in 41.4% of unregulated opioids (72 of 174 samples), whereas fentanyl was detected in 44.8% (78 samples).
- Xylazine detection by FTIR has been increasing in the region. Xylazine was found in 6.9% of unregulated opioid samples in February (12 of 174 samples).
- Drug checking is available in several communities across the Interior Health region. The Medicine Shoppe Pharmacy offers drug checking services in Vernon at #100-3605 31st Street. For updated times and locations in Interior Health, visit https://drugchecking.ca/








BRITISH COLUMBIA CENTRE ON SUBSTANCE USE

www.drugcheckingbc.ca

Drug Checking in Interior Health



PUBLIC HEALTH NOTIFICATIONS


Date & Location	Expected Drug	Drugs Detected	Fentanyl Strip	Benzo Strip	Alert Message
February 14, 2024 Vernon	Down	Fentanyl	Positive	Negative	Substance that looks like meth is sold as down and contains undiluted fentanyl in Vernon.
February 16, 2024 Cranbrook	Down	Fentanyl, Caffeine, Bromazolam, Uncertain Carbohydrate	Positive	Positive	Pinkish-purple chunks sold as down tested positive for high concentrations fentanyl (14%), and bromazolam (10%) in Cranbrook.*

*Quantitative results provided by complementary testing partner <u>Substance</u> using Paper Spray Mass Spectrometry, all other spectroscopy results are determined by FTIR.

Health authorities and community organizations issue further toxic drug alerts from sources other than drug checking. See their respective websites or social media accounts for more alerts.

For information about public health alerts in the Interior Health region, please visit:

Percentage of opioids testing positive for benzodiazepines in the past 6 months




During the month of February, **73.3%** of expected opioid samples tested positive for benzodiazepines using test strips in our partner sites in Interior Health **(137 samples of 187 checked)**. Opioid samples are checked for benzodiazepine-positivity using BTNX test strips and the FTIR spectrometer. The results presented here are derived from both of these technologies and are presumptive until confirmed by a laboratory.

Drug Checking in Interior Health

Number of drugs checks at each site

Location	# of Drugs Checked
ANKORS (Cranbrook)	66
ANKORS (Grand Forks)	24
ANKORS (Nelson)	79
ANKORS (Trail)	9
ASK Wellness (Kamloops)	39
ASK Wellness (Penticton)	51
Medicine Shoppe Pharmacy (Vernon)	36
UBCO HaRT (Kelowna)	118
Total	430

Less than 5 samples were tested at each of the following locations, which are included in the total: ANKORS (Castlegar), ANKORS (Invermere), ASK Wellness (Merritt), ASK Wellness (Williams Lake).




Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

Depressant include: benzodiazepines, etizolam, GHB, hypnotics

Opioid include: "down", heroin, fentanyl (unregulated opioids), and pharmaceutical opioids **Polysubstance includes:** cross-category mixtures Psychedelic include: MDMA and related, 2C-family, tryptamines, ketamine, LSD Stimulant include: methamphetamine, "speed," cocaine and crack cocaine, cathinones Unknown includes: samples where the individual was unable to identify an expected substance - this includes found samples.

Number of opioid samples that matched expectation

using FTIR/test strip drug checking



Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents. 'Down' can refer to any opioid drug present in any amount.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazapine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of <u>both</u> the FTIR <u>and</u> BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners:



February

2024







ASKWELLNESS.CA

