

Drug Checking in Northern Health

Key Findings

- In May, a total of 76 drug checks were performed by services offering FTIR drug checking in the Northern Health region (4 access points).
- The percentage of all opioids testing positive for benzodiazepines by FTIR and/or test strip decreased to 72.4% (42 of 58 samples). The true rate may be higher as benzo-like substances such as etizolam are not reliably detected with benzodiazepine test strips.
- Benzodiazepines were detected by FTIR in 10.7% of unregulated opioids (6 of 56 samples), indicating their presence in higher concentrations.
- Multiple fentanyl analogues have been detected in unregulated opioids. The most recent to emerge, ortho-Methylfentanyl, was detected by FTIR in 1.8% of samples (1 of 56). Fluorofentanyl was detected in 23.2% (13 samples).
- Trends are hard to infer due to the small number of samples over a large region.
- To locate drug checking services in the Northern Health region, please see our site locator.









Drug Checking in Northern Health

Number of samples tested with fentanyl present






To subscribe to the provincial drug alert and response system, text the word JOIN to 253787 (ALERTS).

Number of drugs checks at each site

Location	# of drug checks		
POUNDS (Prince George)	76		

Percentage of opioids testing positive for benzodiazepines in the past 6 months




During the month of May, **72.4**% of expected opioid samples tested positive for benzodiazepines using test strips in our partner site in Northern Health **(42 samples of 58 checked)**. Opioid samples are checked for benzodiazepine-positivity using BTNX test strips and the FTIR spectrometer. The results presented here are derived from both of these technologies and are presumptive until confirmed by a laboratory.

Drug Checking in Northern Health

Number of samples with expected drug present

using FTIR/test strip drug checking



Number of opioid samples with expected drug present

using FTIR/test strip drug checking

EXPECTED DRUG		Present	Not Present	Presence Not Determined	Samples Tested
Heroin		1	0	0	1
Down		55	0	0	55
Pharmaceutical	I	1	0	1	2
Total		57	0	1	58

Please note that the presence of the expected substance does not imply purity, as samples frequently contain adulterating cutting agents

Depressant include: benzodiazepines, etizolam, GHB, hypnotics Opioid include:
"down", heroin, fentanyl
(unregulated opioids), and
pharmaceutical opioids

Polysubstance includes: cross-category mixtures

Psychedelic include: MDMA and related, 2C-family, tryptamines, ketamine, LSD

Stimulant include: methamphetamine, "speed," cocaine and crack cocaine, cathinones **Unknown includes:** Samples where the individual

was unable to identify an expected substance. This includes found samples.

Data represented here are collected from our partner sites across the province. Drug samples are tested using the Fourier Transform Infrared (FTIR) spectrometer in combination with fentanyl test strips and benzodiazapine test strips.

There is 5% fentanyl detection limit on the FTIR spectrometer (McCrae, 2019), and a drug check on any given sample consists of <u>both</u> the FTIR <u>and</u> BTNX fentanyl immunoassay test strip testing done in combination. When applicable, BTNX benzodiazepine immunoassay test strips are also used.

BCCSU gratefully acknowledges the contributions of the following partners:





